

Fractal analysis of ULF electromagnetic emissions in possible association with earthquakes in China

Y. Ida¹⁾, D. Yang²⁾, Q. Li²⁾, H. Sun³⁾, and M. Hayakawa^{1,4,5)}

- 1) *Research Station on Seismo Electromagnetics, The University of Electro-Communications (UEC), 1-5-1 Chofugaoka, Chofu Tokyo 182-8585, JAPAN*
- 2) *Institute of Geophysics, China Earthquake Administration, Beijing 100081, CHINA*
- 3) *Kashi observatory, Earthquake Administration of Xinjiang Uygur Antonomous Region, Xinjiang 844000, CHINA*
- 4) *Hayakawa Institute of Seismo-Electromagnetics, Co. Ltd., UEC Incubation Center, Chofu Tokyo, JAPAN*
- 5) *Earthquake Analysis Laboratory, Information Systems Inc., Kita-aoyama 2-12-42- 305, Minato-ku, Tokyo 107-0061, JAPAN*

The long-term data (during the period of 1 March 2003 through 31 December 2006) of ULF geomagnetic variations observed at Kashi station (geographic coordinates: 39.5°N, 76.0°E) in China, have been used to investigate the long-term variation of fractal dimension of ULF emissions. We have studied the changes in fractal dimension in association with several earthquakes around the observing station. It is then found that a significant change (or decrease) in the fractal dimension of the Z component has taken place before the September 1, 2003 earthquake, which lends a further support to our previous finding based on our improved polarization analysis for the same earthquake. The results obtained are discussed in the contexts of a few aspects (detectability of seismogenic emissions, comparison with previous results by other analysis methods, the importance of fractal analysis in the nonlinear process of the lithosphere and earthquake prediction).